

# Sleep Environment Recommendations for Future Spaceflight Vehicles

Zachary A. Caddick<sup>1</sup>, Kevin Gregory<sup>1</sup>, and Erin E. Flynn-Evans<sup>2</sup>

<sup>1</sup>San Jose State University Research Foundation, San Jose CA, USA and

<sup>2</sup>NASA Ames Research Center, Moffett Field CA, USA {erin.e.flynn-evans@nasa.gov}

**Abstract.** Evidence from spaceflight and ground-based missions demonstrate that sleep loss and circadian desynchronization occur among astronauts, leading to reduced performance and, increased risk of injuries and accidents. We conducted a comprehensive literature review to determine the optimal sleep environment for lighting, temperature, airflow, humidity, comfort, intermittent and erratic sounds, privacy and security in the sleep environment. We reviewed the design and use of sleep environments in a wide range of cohorts including among aquanauts, expeditioners, pilots, military personnel, and ship operators. We also reviewed the specifications and sleep quality data arising from every NASA spaceflight mission, beginning with Gemini. We found that the optimal sleep environment is cool, dark, quiet, and is perceived as safe and private. There are wide individual differences in the preferred sleep environment; therefore modifiable sleeping compartments are necessary to ensure all crewmembers are able to select personalized configurations for optimal sleep.

**Keywords:** Extreme Environments · Habitability · Human Factors · Sleep

## 1 Introduction

Sleep quality -- including the ability to fall asleep and remain asleep -- and sleep duration are dependent upon circadian phase, length of prior wake duration, and time within the sleep episode [1-3]. Proper alignment of scheduled sleep episodes to the circadian pacemaker is important for sleep consolidation and sleep structure [4-5]. High sleep efficiency is best maintained for eight hours when sleep is initiated approximately six hours before the endogenous circadian minimum of core body temperature [4-5]. This phase relationship between the rest-activity cycle and the endogenous circadian timing system implies that even small circadian phase delays of the sleep propensity rhythm with respect to the rest-activity schedule can result in sleep onset insomnia or substantial wake after sleep onset.

In order to quantify the impact of a sub-optimal sleep environment on sleep quality and duration, it is important to measure sleep outcomes when sleep is appropriately timed relative to the circadian and homeostatic drives for sleep. It is possible for an individual to experience sleep disruption in an optimal sleep environment due to the imposed sleep

schedule. Similarly, it is possible for an individual to experience high sleep efficiency in a sub-optimal sleep environment when accumulated sleep debt is present, which dampens the arousal threshold. Our aim was to compile the evidence associated with sleep disruption due to controllable, environmental stimuli in order to aid NASA engineers and operational personnel in the optimal design of crew sleep accommodations for deep spaceflight.

## **2 Methods**

We conducted a comprehensive literature review summarizing optimal sleep hygiene parameters for lighting, temperature, airflow, humidity, comfort, intermittent and erratic sounds, privacy and security in the sleep environment. We reviewed the design and use of sleep environments in a wide range of cohorts including among aquanauts, expeditioners, pilots, military personnel and ship operators. We also reviewed the specifications and sleep quality data arising from every NASA spaceflight mission, beginning with Gemini.

## **3 Recommendations**

The sleep environment required for long duration missions will differ from the sleep accommodations that NASA has developed in the past. Our review revealed several modifications that will be important to make in order to ensure that deep space crews have sleep environments that will provide them with quality sleep.

### **3.1 Sleep Chamber Location**

The location of the sleep station within the vehicle is key to reducing noise and light pollution. Noise emanating from common areas has been shown to be disruptive to sleep [6-7]. Given that there are individual differences in sleep timing preference, it is likely that some crew will chose to be awake, while others are asleep [8-9]. In order to ensure that morning-types and evening-types are both afforded adequate rest, it is desirable to position crew quarters away from the galley area and exercise machinery. We also found that individuals living in a variety isolated and confined environments reported experiencing sleep disruption due to other crewmembers using the waste management system during sleep episodes [9-11]. Therefore, the waste management system should be located far enough away from sleeping quarters that noise is buffered, but close enough that crewmembers are able to quickly access the facility and return to sleep without having to travel too far. It may be appropriate to locate waste management facilities in a module adjacent to the sleep stations.

It is likely that watch schedules will be necessary during deep space missions. We found that in the early history of human spaceflight, watch schedules were very disruptive to sleeping crewmembers due to the close proximity of the sleeping crewmember to the “on watch” crewmember [12]. According to studies of military personnel and pilots,

locating the sleep chambers for off-duty crewmembers away from the command and communication area is desirable [11, 13-15]. However, the sleep chambers should be positioned near enough to the vehicle command center that crewmembers may quickly respond in an emergency situation [11].

### **3.2 Privacy**

It is imperative that each crewmember is provided with a private sleep chamber for the duration of the mission. We found that shared sleep spaces and common bunkrooms are associated with frequent sleep disruption due to other crewmembers [13]. The practice of “hot bunking” has been virtually eliminated from all occupations that we evaluated due to hygiene concerns and the impact that hot bunking has on psychological mood and health [13, 16]. We found that individuals view their sleep location not just as a place for sleep, but also as a space for privacy [7, 14, 16-25]. Access to a private space is viewed as critical to the psychological well-being of individuals living in isolated and confined environments [26]. Similarly, provision for storage of personal items within the sleep chamber was viewed as highly desirable [9, 27]. The sleep chambers for deep space vehicles should also allow crewmembers to customize the space with personal items and reconfiguration of stowage compartments [9].

There have been situations where crewmembers have been displaced from private quarters during spaceflight missions [28]. In these situations it is very difficult for the displaced individuals to obtain adequate sleep [29-31]. Given that the loss of a sleep chamber would likely also be associated with a breach of the spaceflight vehicle, the resulting anxiety may further reduce crewmember sleep quality and quantity. As a result, it is possible that the loss of a sleep chamber could greatly impact the physical and psychological health of crewmembers at a time when successful performance of duties is essential. Given the importance of sleep in conferring fitness for duty, future crew vehicles should include back up, deployable sleep chambers in order to ensure that individuals have access to a private sleep environment throughout the mission.

### **3.3 Habitable Volume**

The crew quarters that are presently on ISS appear to provide enough habitable volume for crewmembers to move as desired during sleep [32-33]. We found one case where a crewmember was too large to fit in the assigned sleep chamber during spaceflight [34]. Although it may be necessary to design all sleep chambers and sleeping bags to the same standard, it is important to consider that larger crewmembers will have less habitable volume relative to smaller crewmembers. As such, it is important to ensure that the crewmembers selected for a deep space mission are able to evaluate the size of the sleep stations in advance of the mission. It may also be desirable to design two sizes for the sleep stations to accommodate larger and smaller crewmembers.

The optimal sleep environment for a planetary excursion will be necessarily different from the optimal sleep environment for spaceflight. During a long duration planetary

excursion, larger crew quarters are necessary due to the comparatively reduced habitable space available in a partial gravity environment. We found that individuals living in isolated and confined environments on Earth use their sleep rooms as a place for privacy and to work in addition to sleep [7, 27, 35]. As a result, the crew rooms on a planetary excursion should include space for a bed (placed horizontally on the floor), a desk and storage of personal belongings. The use of bunkrooms or shared sleep spaces is only appropriate for a short-duration planetary excursion. In these cases, bunks or cots may be used to accommodate crewmembers [7]; however, even during such short excursions private crew quarters would be preferable [27].

### **3.4 Light**

Sleep chambers in spaceflight and on the ground must include features that protect individuals from being awoken by external forces such as light, noise, inadequate temperature and poor air quality. Light is the primary resetting cue for the human circadian pacemaker [36]. Exposure to light at inappropriate times leads to circadian misalignment, which causes sleep disruption [37]. Similarly, exposure to light is alerting and suppresses the drive to sleep [38]. The intensity, spectra, duration, and timing of light determine the magnitude and direction of phase shifting and potency of acute alerting [39]. All wavelengths of light have a negative impact on sleep, but blue light elicits the strongest effect due to the stimulation of intrinsically photosensitive retinal ganglion cells [38]. Exposure to green light is capable of enhancing alertness and suppressing sleep [38], while exposure to red light has the weakest effect on alertness and circadian phase shifting [40]. Evidence from the laboratory, field and subject matter experts support the notion that exposure to light during sleep episodes is disruptive to sleep quality and quantity [12-14, 29, 41-49]. Based on this evidence, all light should be eliminated from the sleep environment. If indicator lights are necessary for identifying egress points, then they should be dim and red [40].

There is strong evidence to suggest that individuals living in isolated and confined environments away from typical solar light dark cues are prone to circadian desynchrony due to self-selecting inappropriate patterns of light exposure [8, 50-54]. This circadian misalignment leads to individuals experiencing a drive to sleep during scheduled wake and an inability to sleep during scheduled sleep opportunities. In order to preserve a stable 24-hour pattern of work and sleep among the crewmembers, it may be desirable to provide a strong cycling of light and darkness in common spaces to mimic the solar light dark cycle and help crewmembers maintain a regular sleep-wake schedule and circadian entrainment [55-56]. However, if such a strategy is utilized, it is important that crewmembers maintain some autonomy in controlling dimmer, personal lighting as would be the case at home on Earth. Similarly, crewmembers scheduled to be on night watch may benefit from supplemental lighting in the vehicle command center in order to enhance alertness and performance [57].

### **3.5 Noise**

Noise is ever-present on space vehicles. We found that noise has been a major cause of sleep disruption throughout the history of spaceflight [12, 19, 29, 58]. The current guidelines allow for exposure to continuous noise above the WHO recommended guidelines [33, 59]. In addition, the current NASA guidelines do not provide mitigations against impulsive or intermittent noise [33]. We found that exposure to intermittent noise is at least as disruptive to sleep as continuous noise exposure [11-12, 15, 19, 29, 58, 60-61]. Given this evidence, exposure to noise be limited to below 35 dB, because exposure to noise above this level is associated with a reduction in sleep quality and quantity, even when individuals do not wake fully [59]. In addition, intermittent noise should be minimized, so that it does not vary beyond 5 dB from background noise levels. There is some evidence to suggest that exposure to continuous white noise less than 25 dB is sufficient to mask intermittent noises [62], therefore it is desirable to allow crewmembers access to white noise in their sleep chamber if desired. Earplugs and/or noise canceling headphones should also be made available for crewmembers [63]. Due to crewmember concerns about missing alarms while wearing earplugs, it may be desirable to develop multi-sensory alarms that include auditory and visual stimulation [64-66].

### **3.6 Temperature and Humidity**

The ambient temperature on early space vehicles varied widely. For optimal sleep, an individual needs to reach his or her thermoneutral equilibrium and should have sufficient bedding available to create a microclimate of between 25-35°C (77-95°F) [67-68]. Given that there are wide individual differences in the optimal temperature for sleep, the sleep environment on future space vehicles should be cool, but there should be sufficient insulation available for crewmembers to modify their environment to suit individual preferences [69-71]. This may mean providing crewmembers with sleeping bags of different thicknesses, or a mechanism for layering sleeping bags together. It is also desirable for sleeping bags to include vents to release heat, because the human core body temperature falls and rises during a typical sleep episode [72]. Warming of proximal and distal skin temperature has been associated with faster sleep onset [73-75] and crewmembers have reported having difficulty sleeping due to cold feet and hands [19, 34], therefore providing a way for crewmembers to warm their extremities prior to sleep may be desirable.

The level of humidity in the environment can also influence sleep quality and quantity. The optimal humidity range for human health is between 40-60% [19]. The presence of humidity in the environment changes the perceived temperature. Higher humidity, with high temperatures are disruptive to sleep [76]. Therefore, lower humidity of 50-60% is optimal for sleep, particularly when ambient temperature is increased.

### **3.7 Air Quality**

The optimal ambient gas mixture for sleep is equivalent to the air experienced at sea level on Earth (78% nitrogen, 21% oxygen, 1% other gases) [16, 21, 77-86]. Similarly, the

optimal air pressure during sleep is equivalent to the pressure on the Earth at sea level [87-88]. Air mixtures that deviate from these conditions, such as what mountaineers experience during expeditions, results in disrupted sleep and periodic breathing [80, 82, 84, 88-90]. In depressurized environments, such as at elevation on Earth, supplemental oxygen can reduce headaches, periodic breathing, and can improve sleep outcomes [91-92]. Airflow is also associated with positive sleep outcomes and aids in the reduction of O<sub>2</sub> [85, 93] and intrusive odors, such as body odor, food, and mechanical smells [12, 34, 85]. Although there is little information on the impact of air pollution and particulates on sleep quality and quantity, reports from lunar expeditions suggest that dust from planetary extra vehicular activities may build up in the habitable environment [29, 34]. As a result, the vents providing airflow to crew sleep chambers should include air filters to protect against crewmembers breathing particulate matter and dust during sleep.

### **3.8 Involuntary Movement**

Involuntary movement due to turbulence is associated with sleep disruption [94]. Therefore, vehicle movement and vibration should be minimized as much as possible. Similarly, the microgravity environment results in the potential for crewmembers to free-float during sleep episodes. Although some crewmembers have reported that they enjoyed that experience, other crewmembers have reported that they prefer to be restrained while sleeping [95]. Given that some individuals may not use harnesses and other attachments, they should be designed, so that they can be removed or secured out of place when not in use. Similarly, separate attachments should be available to secure the sleeping bag to the wall of the sleep chamber if desired.

### **3.9 Summary**

Although we present evidence to support the design of future space vehicles, it is possible that new information will be revealed in the future. NASA supports a great deal of studies in analog and spaceflight environments. As new information becomes available recommendations may evolve and change. Such information should help to further define the optimal sleep environment for deep space transit.

In summary, sleep is critical to crewmember health and performance. In order for crewmembers to achieve optimal sleep, they must be provided with a sleep environment that allows them to achieve quality sleep, free of external disruption. We found that the optimal sleep environment is cool, dark, quiet, and is perceived as safe and private. There are wide individual differences in the preferred sleep environment; therefore modifiable sleeping compartments are necessary to ensure all crewmembers are able to select personalized configurations for optimal sleep. A sub-optimal sleep environment is tolerable for only a limited time, therefore individual sleeping quarters should be designed for long duration missions. In a confined space, the sleep environment serves a dual purpose as a place to sleep, but also as a place for storing personal items and as a place for privacy during non-sleep times. This need for privacy during sleep and wake appears to be

critically important to the psychological well being of crewmembers on long duration missions. Designing sleep chambers for optimal sleep health should produce benefits beyond simply improving sleep quality and quantity on long duration missions.

## References

1. Klerman, E.B., et al., Comparisons of the variability of three markers of the human circadian pacemaker. *J Biol Rhythms*. 17(2), 181--193 (2002)
2. Akerstedt, T. and M. Gillberg, Effects of sleep deprivation on memory and sleep latencies in connection with repeated awakenings from sleep. *Psychophysiology*. 16(1), 49--52 (1979)
3. Wilkinson, R., Some factors influencing the effect of environmental stressors upon performance. *Psychol Bull*. 72(4), 260--272 (1969)
4. Dijk, D.J. and C.A. Czeisler, Paradoxical timing of the circadian rhythm of sleep propensity serves to consolidate sleep and wakefulness in humans. *Neurosci Lett*. 166(1), pp. 63--68 (1994)
5. Dijk, D.J. and C.A. Czeisler, Contribution of the circadian pacemaker and the sleep homeostat to sleep propensity, sleep structure, electroencephalographic slow waves, and sleep spindle activity in humans. *J Neurosci*. 15(5 Pt 1), 3526--3538 (1995)
6. Compton, D.W. and C.D. Benson, *Living and Working in Space: A History of Skylab*. (1983)
7. Hoffman, S.J., Antarctic exploration parallels for future human planetary exploration: A workshop report. National Aeronautics and Space Administration, (2002)
8. Basner, M., et al., Mars 520-d mission simulation reveals protracted crew hypokinesia and alterations of sleep duration and timing. *Proc Natl Acad Sci USA*, 110(7), 2635--2640 (2013)
9. Yan, X.W. and M.E. England, Design evaluation of an arctic research station from a user perspective. *Environment and Behavior*. 33(3), 449--470 (2001)
10. MSFC Skylab Structures, MSFC Skylab structures and mechanical systems mission evaluation, In: NASA TMX-64824, June. National Aeronautics and Space Administration (1974)
11. International Federation of Air Line Pilots' Associations, In-flight flight crew rest facilities. (2013)
12. Hacker, B.C. and J.M. Grimwood, *On the Shoulders of Titans: A History of Project Gemini*. NASA SP-4203. NASA Special Publication, 4203 (1977)
13. Caldwell, J.A., et al., Is fatigue a problem in army aviation: The results of a Survey of aviators and aircrews, US Army Aeromedical Research Laboratory (2000)
14. Green, P.S. Special Report: Sleep Deprivation During Deployment and PTSD. <https://vanwinkles.com/for-our-returning-troops-post-traumatic-sleep-disorders-are-the-new-ptsd>, (2015)
15. Watt, C.G., *Aircraft Fatigue Management*, in Air War College. Air University (2009)
16. Stuster, J.W., *Space station habitability: Recommendations based on a systematic comparative analysis of analogous conditions*. National Aeronautics and Space Administration, (1986)
17. Carrere, S. and G.W. Evans, Life in an Isolated and Confined Environment A Qualitative Study of the Role of the Designed Environment. *Environment and Behavior*, 26(6), 707--741 (1994)
18. Hendrickx, B., The Kamanin Diaries 1969-1971. *Journal-British Interplanetary Society*, 55(9/10), 312--360 (2002)
19. Johnston, R.S., *Skylab Medical Experiments Altitude Test (SMEAT)* (1973)
20. Koros, A., C. Wheelwright, and S. Adam, An evaluation of noise and its effects on shuttle crewmembers during STS-50/USML-1 (1993)

21. MacCallum, T.K. and J. Poynter, Factors affecting human performance in the isolated confined environment of Biosphere 2, In: Third Annual Mid-Atlantic Human Factors Conference, 82--87 (1995)
22. Rosekind, M.R., et al., Crew Factors in Flight Operations XIII: A Survey of Fatigue Factors in Corporate/Executive Aviation Operations, NASA: Moffett Field, CA (2000)
23. Vander Ark, S.T., A.W. Holland, and J. Wood, La Chalupa-30: Lessons Subsea Learned Mission from a 30-day Analogue (1994)
24. Weiss, K., M. Feliot-Rippeault, and R. Gaud, Uses of places and setting preferences in a French Antarctic station. *Environment and Behavior*, 39(2), 147--164 (2007)
25. Willshire, K.F., Human response to vibroacoustic environments of space vehicles. National Aeronautics and Space Administration, Langley Research Center (1984)
26. Harrison, A.A., et al., Implications of privacy needs and interpersonal distancing mechanisms for space station design (1988)
27. Yan, X.W., et al., A critical review of design and use of field tent shelters in polar regions. *Polar Record*, 34(189), 113--122 (1998)
28. Legler, R.D. and F.V. Bennett, Space shuttle missions summary. National Aeronautics and Space Administration: Mission Operations, Johnson Space Center (2011)
29. National Aeronautics and Space Administration, Apollo 11 Mission Report. Houston, TX (1971)
30. National Aeronautics and Space Administration, Apollo 13 Mission Report. Houston, TX (1970)
31. Shepard Jr, A., Apollo 14 mission report (1972)
32. Allen, C. and S. Denham, International Space Station Acoustics - A Status Report. In: 41st International Conference on Environmental Systems (2011)
33. Broyan Jr., J.L., M.A. Borrego, and J.F. Bahr, International Space Station USOS Crew Quarters Development. SAE International (2008)
34. Bluth, B. and M. Helppie, Soviet space stations as analogs (1986)
35. European Space Agency, Mars 500: Isolation Study (2010)
36. Czeisler, C.A. and J.J. Gooley, Sleep and circadian rhythms in humans. *Cold Spring Harb Symp Quant Biol.* 72, 579--597 (2007)
37. Flynn-Evans, E.E., et al., Circadian misalignment affects sleep and medication use before and during spaceflight. *npj Microgravity.* 2, 15019 (2016)
38. Lockley, S.W., et al., Short-wavelength sensitivity for the direct effects of light on alertness, vigilance, and the waking electroencephalogram in humans. *Sleep.* 29(2), 161--168 (2006)
39. Lockley, S.W., Timed melatonin treatment for delayed sleep phase syndrome: the importance of knowing circadian phase. *Sleep.* 28(10), 1214--1216 (2005)
40. Mien, I.H., et al., Effects of exposure to intermittent versus continuous red light on human circadian rhythms, melatonin suppression, and pupillary constriction. *PLoS one.* 9(5), e96532 (2014)
41. Bierman, A., M.G. Figueiro, and M.S. Rea, Measuring and predicting eyelid spectral transmittance. *Journal of Biomedical Optics.* 16(6), 067011--067011 (2011)
42. Cho, J.R., et al., Let there be no light: the effect of bedside light on sleep quality and background electroencephalographic rhythms. *Sleep medicine.* 14(12), 1422--1425 (2013)
43. Dijk, D.-J., et al., Sleep, circadian rhythms, and performance during space shuttle missions. *The Neurolab Spacelab Mission: Neuroscience Research in Space.* 211--221 (2003)
44. Figueiro, M.G. and M.S. Rea, Short-wavelength light enhances cortisol awakening response in sleep-restricted adolescents. *International Journal of Endocrinology.* 2012 (2012)

45. Figueiro, M.G., B. Plitnick, and M.S. Rea, Pulsing blue light through closed eyelids: Effects on acute melatonin suppression and phase shifting of dim light melatonin onset. *Nature and Science of Sleep*. 6, 149 (2014)
46. Grandner, M.A., et al., Short wavelength light administered just prior to waking: A pilot study. *Biological Rhythm Research*. 44(1), 13--32 (2013)
47. Potter, J.J., et al., Polar field tent shelters and well-being of users. *Environment and behavior*. 30(3), 398--420 (1998)
48. Thompson, A., et al., Effects of dawn simulation on markers of sleep inertia and post-waking performance in humans. *Eur J Appl Physiol*. 114(5), 1049--1056 (2014)
49. Zeitzer, J.M., et al., Millisecond flashes of light phase delay the human circadian clock during sleep. *J Biol Rhythms*. 29(5), 370-376 (2014)
50. Arendt, J., Biological rhythms during residence in polar regions. *Chronobiology International*. 29(4), 379-394 (2012)
51. Halberg, F., et al., Human biological rhythms during and after several months of isolation underground in natural caves. *Bulletin of The National Speleological Society*. 32, 89-115 (1970)
52. Kleitman, N., Sleep and Wakefulness. *The American Journal of the Medical Sciences*. 249(2), 140 (1965)
53. Miller, N.L. and J. Nguyen, Working the nightshift on the USS John C. Stennis: Implications for enhancing warfighter effectiveness. In: *Proceedings of the Human-Systems Integration Symposium* (2003)
54. Siffre, M., Biological rhythms, sleep, and wakefulness in prolonged confinement, in *ESA, Proceedings of the Colloquium on Space and Sea*. 53--68 (1988)
55. Duplessis, C.A., et al., Submarine watch schedules: Underway evaluation of rotating (contemporary) and compressed (alternative) schedules. *Undersea & Hyperbaric Medicine*. 34, 21--33 (2007)
56. Young, C.R., et al., At-sea trial of 24-h-based submarine watchstanding schedules with high and low correlated color temperature light sources. *Journal of Biological Rhythms*. 30, 144--154 (2015)
57. Barger, L.K., et al., Sleep and cognitive function of crewmembers and mission controllers working 24-h shifts during a simulated 105-day spaceflight mission. *Acta Astronautica*. 93, pp. 230--242 (2014)
58. Flynn-Evans, E.E., et al. Sleep duration, disruption and hypnotic use among 76 astronauts on short duration missions. in *NASA Human Research Program Meeting*. Houston, TX (2012)
59. World Health Organization, *Night noise guidelines for Europe* (2009)
60. Fyhri, A. and G.M. Aasvang, Noise, sleep and poor health: Modeling the relationship between road traffic noise and cardiovascular problems. *Science of the Total Environment*. 408, 4935--4942 (2010)
61. Schmidt, F.P., et al., Effect of nighttime aircraft noise exposure on endothelial function and stress hormone release in healthy adults. *European Heart Journal*. eht269 (2013)
62. Stanchina, M.L., et al., The influence of white noise on sleep in subjects exposed to ICU noise. *Sleep Med*. 6(5), 423--428 (2005)
63. Department of the Army, *Manual 6-22.5: Combat and Operational Stress Control Manual for Leaders and Soldiers*. Washington, DC: US Dept. of the Army. 111 (2009)
64. Kawada, T. and S. Suzuki, Change in rapid eye movement (REM) sleep in response to exposure to all-night noise and transient noise. *Archives of Environmental Health*. 54, 336--340 (1999)
65. Muzet, A., Environmental noise, sleep and health. *Sleep Medicine Reviews*. 11, 135--142 (2007)

66. Rechtschaffen, A., P. Hauri, and M. Zeitlin, Auditory awakening thresholds in REM and NREM sleep stages. *Perceptual and Motor Skills*. 22(3), 927-942 (1966)
67. Okamoto, K., et al., A survey of bedroom and bed climate of the elderly in a nursing home. *Applied Human Science*. 17(3), 115--120 (1998)
68. Okamoto-Mizuno, K., K. Tsuzuki, and K. Mizuno, Effects of head cooling on human sleep stages and body temperature. *International Journal of Biometeorology*. 48, 98--102 (2003)
69. Haskell, E.H., et al., The effects of high and low ambient temperatures on human sleep stages. *Electroencephalography and Clinical Neurophysiology*. 51, 494-501 (1981)
70. Lin, Z. and S. Deng, A study on the thermal comfort in sleeping environments in the subtropics — Measuring the total insulation values for the bedding systems commonly used in the subtropics. *Building and Environment*. 43, 905--916 (2008)
71. Häuplik-Meusburger, S., *Architecture for Astronauts: An Activity-based Approach*. Springer Science & Business Media (2011)
72. Van Someren, E.J.W., et al., Circadian and age-related modulation of thermoreception and temperature regulation: Mechanisms and functional implications. *Ageing Research Reviews*. 1, 721--778 (2002)
73. Kräuchi, K., et al., Warm feet promote the rapid onset of sleep. *Nature*. 401, 36--37 (1999)
74. Raymann, R.J.E.M., D.F. Swaab, and E.J.W. Van Someren, Cutaneous warming promotes sleep onset. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*. 288, R1589--R1597 (2005)
75. Raymann, R.J.E.M., D.F. Swaab, and E.J.W. Van Someren, Skin deep: Enhanced sleep depth by cutaneous temperature manipulation. *Brain*. 131(2), 500--513 (2008)
76. Okamoto-Mizuno, K., et al., Effects of partial humid heat exposure during different segments of sleep on human sleep stages and body temperature. *Physiology & Behavior*. 83, 759--765 (2005)
77. Daues, K.R., *A history of spacecraft environmental control and life support systems* (2006)
78. Gothe, B., et al., Effect of quiet sleep on resting and CO<sub>2</sub>-stimulated breathing in humans. *Journal of Applied Physiology*. 50(4), 724--730 (1981)
79. Lo, Y.-L., et al., Genioglossal muscle response to CO<sub>2</sub> stimulation during NREM sleep. *Sleep*. 29(4), 470 (2006)
80. Reite, M., et al., Sleep physiology at high altitude. *Electroencephalography and Clinical Neurophysiology*. 38(5), 463--471 (1975)
81. Robin, E.D., et al., Alveolar gas tensions, pulmonary ventilation and blood pH during physiologic sleep in normal subjects. *Journal of Clinical Investigation*, 37(7), 981 (1958)
82. Salvaggio, A., et al., Effects of high-altitude periodic breathing on sleep and arterial oxyhaemoglobin saturation. *European Respiratory Journal*. 12(2), 408--413 (1998)
83. Schiffman, P., et al., Sleep deprivation decreases ventilatory response to CO<sub>2</sub> but not load compensation. *CHEST Journal*. 84(6), 695--698 (1983)
84. Selvamurthy, W., et al., Sleep patterns at an altitude of 3500 metres. *International Journal of Biometeorology*. 30, 123--135 (1986)
85. Strøm-Tejsen, P., et al., The effects of bedroom air quality on sleep and next-day performance. *Indoor Air*. (2015)
86. Szymusiak, R. and E. Satinoff, Maximal REM sleep time defines a narrower thermoneutral zone than does minimal metabolic rate. *Physiology & Behavior*. 26, 687--690 (1981)
87. Miller, J.C. and S. Horvath, Sleep at altitude. *Aviation, space, and environmental medicine*. 48(7), 615--620 (1977)
88. Mizuno, K., K. Asano, and N. Okudaira, Sleep and respiration under acute hypobaric hypoxia. *The Japanese Journal of Physiology*. 43, 161--175 (1993)

89. Dietz, T.E. An Altitude Tutorial. [http://www.ismmed.org/np\\_altitude\\_tutorial.htm](http://www.ismmed.org/np_altitude_tutorial.htm) (2006)
90. Muza, S.R., C.S. Fulco, and A. Cymerman, Altitude Acclimatization Guide. Army Research Inst of Environmental Medicine Natick MA Thermal and Mountain Medicine Division (2004)
91. Barash, I.A., et al., Nocturnal oxygen enrichment of room air at 3800 meter altitude improves sleep architecture. *High Altitude Medicine & Biology*. 2(4), 525--533 (2001)
92. Luks, A.M., et al., Room oxygen enrichment improves sleep and subsequent day-time performance at high altitude. *Respiration physiology*. 113(3), 247--258 (1998)
93. Federal Aviation Administration, Flightcrew Member Duty and Rest Requirements (2011)
94. Matsangas, P., N.L. Shattuck, and M.E. McCauley, Sleep duration in rough sea conditions. *Aerospace Medicine and Human Performance*. 86, 901--906 (2015)
95. Petty, J.I. Space Shuttle Life, <http://spaceflight.nasa.gov/history/shuttle-mir/history/h-b-sslife.htm> (2013)